

Alternative products to treat allergic rhinitis and alternative routes for allergy immunotherapy

Kagan Ipci, M.D.,¹ Tugba Oktemer, M.D.,² Nuray Bayar Muluk, M.D.,³ Ethem Şahin, M.D.,⁴ Niyazi Altıntoprak, M.D.,⁵ Sameer Ali Bafaqeh, M.D.,⁶ Yasemin Kurt, M.D.,⁷ Ranko Mladina, M.D.,⁸ Marin Šubarić, M.D.,⁹ and Cemal Cingi, M.D.¹⁰

ABSTRACT

Background: Some alternative products instead of immunotherapy are used in patients with allergic rhinitis (AR).

Methods: In this paper, alternative products to treat allergic rhinitis and alternative routes for allergy immunotherapy are reviewed.

Results: Alternative products and methods used instead of immunotherapy are tea therapy, acupuncture, *Nigella sativa*, cinnamon bark, Spanish needle, acerola, capsaicin (*Capsicum annuum*), allergen-absorbing ointment, and cellulose powder. *N. sativa* has been used in AR treatment due to its anti-inflammatory effects. *N. sativa* oil also inhibits the cyclooxygenase and 5-lipoxygenase pathways of arachidonic acid metabolism. The beneficial effects of *N. sativa* seed supplementation on the symptoms of AR may be due to its antihistaminic properties. To improve the efficacy of immunotherapy, some measures are taken regarding known immunotherapy applications and alternative routes of intralymphatic immunotherapy and epicutaneous immunotherapy are used.

Conclusion: There are alternative routes and products to improve the efficacy of immunotherapy.

(Am J Rhinol Allergy 30, S8–S10, 2016; doi: 10.2500/ajra.2016.30.4364)

Allergen immunotherapy (AIT) works by stimulating dendritic cells (DC), T cells, and B cells. By contrast, adverse effects are mediated by mast cells and blood vessels. The ideal route for AIT is characterized by a high density of DCs, T cells, and B cells but a low density of mast cells and blood vessels. Lymph nodes contain masses of DCs, T cells, and B cells but few mast cells. Intralymphatic AIT (ILIT) was shown to be safe, and the number of allergen injections could be reduced to three. The epidermis contains dense DCs and no mast cells or blood vessels. Epicutaneous AIT (EPIT) was also shown to be safe, and the allergen administrations could be reduced to six patches.¹

The number of allergen administrations can be reduced by enhancing the immunologic effects of each administration. AIT works by inducing blocking antibodies, T-helper 1 and regulatory T cells, orchestrated by DCs. Immunologic effects are classically enhanced by adjuvants. Replacing alum with bacterial products, such as the lipopolysaccharide derivative monophosphoryl lipid A or bacterial DNA, allows for a reduction in the number of injections to four or six.^{1,2–4}

Alternative medicine refers to nontraditional medical care used instead of mainstream medical care. Some alternative products are used in patients with AR. Alternative products and methods used for treatment of AR are tea therapy, acupuncture, and *Nigella sativa* Linn. (Ranunculaceae), Cinnamon bark, Spanish needle, and acerola; capsaicin (*Capsicum annuum*), allergen-absorbing ointment and cellulose. Alternative routes of immunotherapy have been studied, including

ILIT and EPIT. This article reviewed novel alternative routes of allergy immunotherapy administration as well as the use of alternative medicine products for the treatment of allergic diseases.

ALTERNATIVE ROUTES

ILIT

In lymph nodes, the density of DCs, B cells, and T cells is maximal, whereas the density of mast cells is low. Direct injection into a subcutaneous lymph node has been shown to be safe and to ameliorate symptoms of AR after three injections.¹ ILIT aims to improve treatment efficacy and safety, and to reduce treatment duration, which, thereby, makes allergen immunotherapy more patient friendly. In different studies, ILIT was administered three times at 4-week intervals.^{5,6} ILIT with grass-pollen or birch-pollen extracts seems to reduce nasal allergic symptoms. Senti *et al.*⁷ showed that ILIT was safe and induced allergen tolerance after three injections in a randomized double-blind trial ILIT with Modular-Antigen-Translocation-Feld 1 (cat antigen) in alum. Hence, ILIT might constitute a less time-consuming and more cost-effective alternative to conventional subcutaneous allergen-specific immunotherapy.⁶ Through injection into a lymph node, it is hoped that ILIT will enhance the development of protective immunity.^{8,9} Clinical trials conducted to date demonstrated that ILIT may be clinically effective after only a few injections and induces allergen-specific immunoglobulin G as demonstrated with subcutaneous immunotherapy.^{10,11}

EPIT

The epidermis is characterized by a high density of potent DCs and an absence of blood vessels. Therefore, AIT administered *via* this route should be more efficient and safer. EPIT has been demonstrated to be safe and to ameliorate the symptoms of AR after six allergen patch applications.¹ In EPIT, an allergen is administered to the non-vascularized epidermis.¹² However, as an advantage of EPIT over sublingual immunotherapy, keratinocytes can additionally be activated by physical irritation, *e.g.*, abrasion or adhesive tape stripping, or by adding adjuvants.¹³ Epithelial damage increases the keratinocyte expression of additional molecules of interleukin (IL) 1 α , IL-6, and tumor necrosis factor α , skewing the immune response toward a [T-helper type 1] type response.¹⁴ Such activation of keratinocytes is important to create a proinflammatory environment with enhanced activation of Langerhans cells.

From the ¹Ear, Nose and Throat (ENT) Department, Ankara Koru Hospital, Ankara, Turkey, ²ENT Department, Private Polatlı Can Hospital, Polatlı-Ankara, Turkey, ³ENT Department, Faculty of Medicine, Kırıkkale University, Kırıkkale, Turkey, ⁴ENT Clinics, Bayındır İçerenköy Hospital, Istanbul, Turkey, ⁵ENT Clinics, Tuzla State Hospital, Istanbul, Turkey, ⁶ENT Department, King Saud University, Riyadh, Saudi Arabia, ⁷Family Medicine Department, Eskisehir Osmangazi University, Eskisehir, Turkey, ⁸ENT Department, University Hospital, Rebro-KBC Zagreb, Croatia, ⁹ENT Department, Clinical Hospital Dubrava, Zagreb, Croatia, and ¹⁰ENT Department, Faculty of Medicine, Eskisehir Osmangazi University, Eskisehir, Turkey
Preparation of this manuscript, including design and planning, was supported by Continuous Education and Scientific Research Association. The authors extend their appreciation to the International Scientific Partnership Program (ISPP) at King Saud University for funding this research work through ISPP# 0041
The authors have no conflicts of interest to declare pertaining to this article
Address correspondence to Nuray Bayar Muluk, M.D., Birlik Mahallesi, Zirvekent 2, Etap Sitesi, C-3 blok, no. 62/43, 06610 Çankaya, Ankara, Turkey
E-mail address: nuray.bayar@yahoo.com, nurayb@hotmail.com
Copyright © 2016, OceanSide Publications, Inc., U.S.A.

Besides enhancing the penetration of allergens by removing the stratum corneum,¹⁵ "repeated tape stripping also functions as a 'physical' adjuvant through the activation of keratinocytes, which then secrete various proinflammatory cytokines (IL-1, IL-6, IL-8, [tumor necrosis factor] α and [interferon] γ), which favor the maturation and emigration of DCs to the draining lymph nodes."^{16,17} The results of the first pilot trial revealed that patients treated with 12 patches that contained pollen extract showed significant alleviation of hay fever symptoms compared with patients who received placebo. In line with the historic study results described above, no severe systemic allergic reactions were reported. The only adverse events observed were mild local eczematous reactions under the skin patch.¹⁸

EPIT holds promise in all four aspects: (1) the epidermis contains a high number of potent antigen presenting cells, (2) adjuvants can be topically administered and/or physical or chemical trauma to keratinocytes may already act as a physical adjuvant, (3) the epidermis is nonvascularized and it decreases systemic exposure to allergen and thus decreases the risk of a systemic reaction, and (4) epicutaneous administration can be performed at home and is painless.¹²

ALTERNATIVE PRODUCTS AND METHODS

Tea Therapy

In Asia, tea therapy is an easy-to-administer self-care method for patients with AR. Japanese green tea with methylated catechin as an active ingredient is thought to be beneficial for patients with AR who are sensitized to cedar and to relieve symptoms, improve quality of life, and reduce the peripheral eosinophil count.¹⁹ Benifuuki green tea contains O-methylated epigallocatechin-3-O-(3-O-methyl) gallate, a compound that has antiallergic properties.²⁰⁻²²

Acupuncture

Acupuncture can be recommended as an adjunctive therapy for AR. Acupuncture and acupressure work *via* stimulating specific points on the human body.¹⁹ Based on the latest meta-analysis of studies published from 1980 to 2013, acupuncture seems to be a safe, valid, and cost-effective option for allergy relief, and can be recommended as an adjunctive therapy for AR.²³ Acupuncture decreases nasal symptom scores and improves quality of life with no serious systemic reaction or adverse effects. Its efficiency and safety heavily depend on the clinical experience and skill of the acupuncturist.^{19,23} Zhang *et al.*²⁴ investigated the efficacy and safety of ear acupressure (EAP), which is a noninvasive semi-self-administered form of acupuncture, in perennial AR (PAR). Participants with PAR were randomized to receive real or sham EAP treatment once a week for 8 weeks and then were followed-up for 12 weeks. The participants were instructed to administer EAP stimulation three times daily. They concluded that EAP showed short-term and extended benefit for improving PAR symptoms and quality of life for patients with PAR.²⁴

Herbal Products for AR

N. sativa. The seeds of *N. sativa*, commonly known as black seed or black cumin, are used in folk (herbal) medicine throughout the world for the treatment and prevention of several diseases and conditions, which include asthma, diarrhea, and dyslipidemia.²⁵ The seeds contain fixed and essential oils, proteins, alkaloids and saponin. Much of the biologic activity of the seeds has been shown to be due to thymoquinone, the major component of the essential oil, but the activity is also present in the fixed oil.²⁵ *N. sativa* has antiallergic, antioxidant, and anti-inflammatory effects on patients with allergy. Furthermore, *N. sativa* shows beneficial effects on asthma.²⁶ Boskabady *et al.*²⁷ reported the antiasthmatic (bronchodilatory) effect of boiled extract of *N. sativa* in the airways of patients with asthma. *N. sativa* (black seed)

is capable of relieving the symptoms of allergic diseases (AR, atopic eczema, bronchial asthma).²⁸

Effects of N. sativa on AR. In the prospective, double-blind clinical trial of patients with AR by Nikakhlagh *et al.*²⁹ *N. sativa* was reported to reduce the presence of nasal mucosal congestion, nasal itching, runny nose, sneezing attacks, turbinate hypertrophy, and mucosal pallor during the first 2 weeks. In a study by Kalus *et al.*²⁸ they reported that black seed oil was an effective adjuvant for the treatment of allergic diseases (AR, bronchial asthma, atopic eczema). Işık *et al.*³⁰ reported that *N. sativa* seed supplementation during specific immunotherapy for AR may be considered a potential adjuvant therapy. The beneficial effects of *N. sativa* seed supplementation on the symptoms of AR may be due to its antihistaminic properties.³⁰ Chakravarty³¹ demonstrated that, when rat macrophages were incubated with nigellone thymoquinone (a carbon polymer) obtained from *N. sativa* seed, nigellone inhibited histamine release from macrophages.

Cinnamon Bark, Spanish Needle, and Acerola. The botanical product that is a combination of cinnamon bark extract, dehydrated Spanish needle (*Bidens pilosa*) leaf and stem, and acerola fruit concentrate inhibits prostaglandin D₂ and shows a steroid-like effect. It decreases allergic inflammation in the late phase.^{32,33}

Capsaicin. Capsaicin (*C. annuum*), derived from red peppers, may work by desensitizing nasal nerve fibers and reducing nasal hyperresponsiveness.^{32,34} It helps to improve nasal congestion and is commercially available.³⁵

Allergen-Absorbing Ointment. Petroleum-based ointment that contains long-chain hydrocarbons can block allergen absorption into the nasal mucosa. When applied three times daily for 14 days, nasal symptom scores improved in both adults and children.^{33,36-38}

Cellulose Powder. Inert and micronized cellulose powder has been speculated to block mucosal allergen absorption after intranasal application.³³

Other Herbal Preparations. There are some other herbal preparations that have shown limited efficacy or a lack of efficacy in the treatment of AR and/or conjunctivitis. These are quercetin (bioflavonoids),³⁹ stinging nettle (*Urtica dioica*) (radix urticae),⁴⁰ *Perilla frutescens* (an Asian herb),⁴¹ Eucalyptus (generates an oil rich in 1,8-cineole),⁴² and grape seed extract.⁴³

CONCLUSION

REFERENCES

1. Senti G, and Kündig TM. AIT for allergic rhinitis—New delivery options. In Global Atlas of Allergic Rhinitis and Chronic Rhinosinusitis. Akdis CA, Hellings P, and Agache I (Eds). European Academy of Allergy and Clinical Immunology, Zurich, Switzerland, 217–218, 2015.
2. Warshakoon HJ, Hood JD, Kimbrell MR, et al. Potential adjuvant properties of innate immune stimuli. Hum Vaccin 5:381–394, 2009.
3. Wang Q, Zhou Z, Tang S, and Guo Z. Carbohydrate-monophosphoryl lipid A conjugates are fully synthetic self-adjuvanting cancer vaccines eliciting robust immune responses in the mouse. ACS Chem Biol 7:235–240, 2012.
4. Mata-Haro V, Cekic C, Martin M, et al. The vaccine adjuvant monophosphoryl lipid A as a TRIF-biased agonist of TLR4. Science 316: 1628–1632, 2007.
5. Senti G, Prinz Vavricka BM, Erdmann I, et al. Intralymphatic allergen administration renders specific immunotherapy faster and safer: A randomized controlled trial. Proc Natl Acad Sci U S A 105:17908–17912, 2008.
6. Hylander T, Latif L, Petersson-Westin U, and Cardell LO. Intralymphatic allergen-specific immunotherapy: An effective and safe alternative treatment route for pollen-induced allergic rhinitis. J Allergy Clin Immunol 131:412–420, 2013.
7. Senti G, Cramer R, Kuster D, et al. Intralymphatic immunotherapy for cat allergy induces tolerance after only 3 injections. J Allergy Clin Immunol 129:1290–1296, 2012.

8. Senti G, Johansen P, and Kundig TM. Intralymphatic immunotherapy: From the rationale to human applications. *Curr Top Microbiol Immunol* 352:71–84, 2011.
9. Valenta R, Campana R, Marth K, and van Hage M. Allergen-specific immunotherapy: From therapeutic vaccines to prophylactic approaches. *J Intern Med* 272:144–157, 2012.
10. Senti G, Prinz Vavricka BM, Erdmann I, et al. Intralymphatic allergen administration renders specific immunotherapy faster and safer: A randomized controlled trial. *Proc Natl Acad Sci U S A* 105:17908–17912, 2008.
11. Senti G, Cramer R, Kuster D, et al. Intralymphatic immunotherapy for cat allergy induces tolerance after only 3 injections. *J Allergy Clin Immunol* 129:1290–1296, 2012.
12. Senti G, von Moos S, and Kundig TM. Epicutaneous immunotherapy for aeroallergen and food allergy. *Curr Treat Options Allergy* 1:68–78, 2014.
13. von Moos S, Johansen P, Waeckerle-Men Y, et al. The contact sensitizer diphenylcyclopropanone has adjuvant properties in mice and potential application in epicutaneous immunotherapy. *Allergy* 67: 638–646, 2012.
14. Swamy M, Jamora C, Havran W, and Hayday A. Epithelial decision makers: In search of the ‘epimune.’ *Nat Immunol* 11:656–665, 2010.
15. Dickel H, Goulioumis A, Gambichler T, et al. Standardized tape stripping: A practical and reproducible protocol to uniformly reduce the stratum corneum. *Skin Pharmacol Physiol* 23:259–265, 2010.
16. Nickoloff BJ, and Naidu Y. Perturbation of epidermal barrier function correlates with initiation of cytokine cascade in human skin. *J Am Acad Dermatol* 30:535–546, 2010.
17. Dickel H, Gambichler T, Kamphowe J, et al. Standardized tape stripping prior to patch testing induces upregulation of Hsp90, Hsp70, IL-33, TNF-alpha and IL-8/CXCL8 mRNA: New insights into the involvement of ‘alarmins’. *Contact Dermatitis* 63:215–222, 2010.
18. Senti G, Graf N, Haug S, et al. Epicutaneous allergen administration as a novel method of allergen-specific immunotherapy. *J Allergy Clin Immunol* 124:997–1002, 2009.
19. Zhang W. Complementary and alternative medicine for allergic rhinitis. In *Global Atlas of Allergic Rhinitis and Chronic Rhinosinusitis*. Akdis CA, Hellings P, and Agache I (Eds). European Academy of Allergy and Clinical Immunology, 228–229, 2015.
20. Fujimura Y, Tachibana H, Maeda-Yamamoto M, et al. Antiallergic tea catechin, (-)-epigallocatechin-3-O-(3-O-methyl)-gallate, suppresses FcεpsilonRI expression in human basophilic KU812 cells. *J Agric Food Chem* 50:5729–5734, 2002.
21. Maeda-Yamamoto M, Inagaki N, Kitaura J, et al. O-methylated catechins from tea leaves inhibit multiple protein kinases in mast cells. *J Immunol* 172:4486–4492, 2004.
22. Maeda-Yamamoto M, Ema K, Monobe M, et al. Epicatechin-3-O-(3'-O-methyl)-gallate content in various tea cultivars (*Camellia sinensis* L.) and its in vitro inhibitory effect on histamine release. *J Agric Food Chem* 60:2165–2170, 2012.
23. Feng S, Han M, Fan Y, et al. Acupuncture for the treatment of allergic rhinitis: A systematic review and meta-analysis. *Am J Rhinol Allergy* 29:57–62, 2015.
24. Zhang CS, Xia J, Zhang AL, et al. Ear acupressure for perennial allergic rhinitis: A multicenter randomized controlled trial. *Am J Rhinol Allergy* 28:e152–e157, 2014.
25. Ali BH, and Blunden G. Pharmacological and toxicological properties of *Nigella sativa*. *Phytother Res* 17:299–305, 2003.
26. Büyüköztürk S, Gelincik A, Özşeker F, et al. *Nigella sativa* (black seed) oil does not affect the T-helper 1 and T-helper 2 type cytokine production from splenic mononuclear cells in allergen sensitized mice. *J Ethnopharmacol* 100:295–298, 2005.
27. Boskabady MH, Mohsenpoor N, and Takaloo L. Antiasthmatic effect of *Nigella sativa* in airways of asthmatic patients. *Phytomedicine* 17:707–713, 2010.
28. Kalus U, Pruss A, Bystron J, et al. Effect of *Nigella sativa* (black seed) on subjective feeling in patients with allergic diseases. *Phytother Res* 17:1209–1214, 2003.
29. Nikakhlagh S, Rahim F, Aryani FH, et al. Herbal treatment of allergic rhinitis: The use of *Nigella sativa*. *Am J Otolaryngol* 32:402–407, 2011.
30. Işık H, Cevikbaş A, Güner US, et al. Potential adjuvant effects of *Nigella sativa* seeds to improve specific immunotherapy in allergic rhinitis patients. *Med Princ Pract* 19:206–211, 2010.
31. Chakravarty N. Inhibition of histamine release from mast cells by nigellone. *Ann Allergy* 70:237–242, 1993.
32. Bielory L. Complementary and alternative therapies for allergic rhinitis and conjunctivitis. In: *Up to Date*. Corren J, and Feldweg AM (Eds); accessed August 2, 2016.
33. Corren J, Lemay M, Lin Y, et al. Clinical and biochemical effects of a combination botanical product (ClearGuard) for allergy: A pilot randomized double-blind placebo-controlled trial. *Nutr J* 7:20, 2008.
34. Blom HM, Van Rijswijk JB, Garrelds IM, et al. Intranasal capsaicin is efficacious in non-allergic, non-infectious perennial rhinitis. A placebo-controlled study. *Clin Exp Allergy* 27:796–801, 1997.
35. Bernstein JA, Davis BP, Picard JK, et al. A randomized, double-blind, parallel trial comparing capsaicin nasal spray with placebo in subjects with a significant component of nonallergic rhinitis. *Ann Allergy Asthma Immunol* 107:171–178, 2011.
36. Bufe A. A simple advice for the prevention of pollen-induced allergic rhinitis. *Int Arch Allergy Immunol* 121:85–86, 2000.
37. Schwetz S, Olze H, Melchisedech S, et al. Efficacy of pollen blocker cream in the treatment of allergic rhinitis. *Arch Otolaryngol Head Neck Surg* 130:979–984, 2004.
38. Geithoff UW, Blum A, Rupp-Classen M, and Plinkert PK. Lipid-based nose ointment for allergic rhinitis. *Otolaryngol Head Neck Surg* 133:754–761, 2005.
39. Remberg P, Björk L, Hedner T, and Sterner O. Characteristics, clinical effect profile and tolerability of a nasal spray preparation of *Artemisia abrotanum* L. for allergic rhinitis. *Phytomedicine* 11:36–42, 2004.
40. Mittman P. Randomized, double-blind study of freeze-dried *Urtica dioica* in the treatment of allergic rhinitis. *Planta Med* 56:44–47, 1990.
41. Takano H, Osakabe N, Sanbongi C, et al. Extract of *Perilla frutescens* enriched for rosmarinic acid, a polyphenolic phytochemical, inhibits seasonal allergic rhinoconjunctivitis in humans. *Exp Biol Med* (Maywood) 229:247–254, 2004.
42. Burrow A, Eccles R, and Jones AS. The effects of camphor, eucalyptus and menthol vapour on nasal resistance to airflow and nasal sensation. *Acta Otolaryngol* 96:157–161, 1983.
43. Bernstein DI, Bernstein CK, Deng C, et al. Evaluation of the clinical efficacy and safety of grape seed extract in the treatment of fall seasonal allergic rhinitis: A pilot study. *Ann Allergy Asthma Immunol* 88:272–278, 2002. □